

FACTORWISE RIGIDITY OF PRODUCTS OF PSEUDO-ARCS

David J. BELLAMY*

Department of Mathematics, University of Delaware, Newark, DE 19716, USA

Judy A. KENNEDY

Tulane University of Louisiana, New Orleans, LA 70118; and, Auburn University, Auburn, AL 36849, USA

Received 25 March 1985

It is proven that every homeomorphism of a product of pseudo-arcs is a composition of a product homeomorphism with a homeomorphism which only permutes the factors.

AMS (MOS) Subj. Class.: Primary 54F20; Secondary 54C20
54B10, 54H15, 54F15

product continuum	ε -mapping
product homeomorphism	ε -homeomorphism
homogeneous continuum	pseudo-arc

A *continuum* is a compact, connected Hausdorff space. The interval $[0, 1]$ is denoted by I . A topological space X is *homogeneous* if and only if for each $x, y \in X$ there is a homeomorphism $h: X \rightarrow X$ such that $h(x) = y$. *Map* or *mapping* always means continuous function.

If X is a continuum then $\mathcal{H}(X)$, the set of homeomorphisms of X is a topological group which acts continuously on X when $\mathcal{H}(X)$ is given the compact open topology [1]. If Γ is an indexing set, and for each $\gamma \in \Gamma$, X_γ is a continuum, then $\prod_{\gamma \in \Gamma} X_\gamma$ is a *product continuum*. In a product continuum $\prod_{\gamma \in \Gamma} X_\gamma$, a *product homeomorphism* is a homeomorphism of the form $H(\langle x_\gamma \rangle_{\gamma \in \Gamma}) = \langle h_\gamma(x_\gamma) \rangle_{\gamma \in \Gamma}$ where each h_γ is a homeomorphism of X_γ . *Product mapping* is defined similarly. If all the X_γ 's are equal to X , $\prod_{\gamma \in \Gamma} X_\gamma$ may be written as X^Γ . If n is a positive integer, I^n is the boundary of I^n ; that is, the set of those points of I^n with at least one coordinate equal to zero or one.

If X and Y are metric continua and $\varepsilon > 0$, an ε -map $f: X \rightarrow Y$ is a map such that for each $y \in Y$, $f^{-1}(y)$ has diameter less than ε . If X is metric, an ε -homeomorphism $h: X \rightarrow X$ is a homeomorphism such that, for each $p \in X$, the distance from p to $h(p)$ is less than ε . The set of ε homeomorphisms is a neighborhood of the identity in $\mathcal{H}(X)$. If each of X_i , $1 \leq i \leq n$, is a metric continuum with metric d_i , the metric

* The first author acknowledges support by NSF grant number MCS 8302176.

d on $\prod_{i=1}^n X_i$ is always taken to be the maximum metric; that is, $d(\langle x_i \rangle_{i=1}^n, \langle y_i \rangle_{i=1}^n)$ is the largest element of the set $\{d_i(x_i, y_i)\}_{i=1}^n$. With this convention, it follows that a product mapping is an ε -map if and only if each of the factor maps is an ε -map, and a product homeomorphism is an ε -homeomorphism if and only if each of its factor maps is an ε -homeomorphism.

A map $f: X \rightarrow I^n$ is *essential* provided there does not exist a nonsurjective mapping $g: X \rightarrow I^n$ such that $g|_{f^{-1}(\dot{I}^n)} = f|_{f^{-1}(\dot{I}^n)}$. If X and Y are continua, a mapping $f: X \rightarrow Y$ is *weakly confluent* if and only if, for every subcontinuum $M \subseteq Y$, there exists a subcontinuum $W \subseteq X$ such that $f(W) = M$.

Lemma 1. *Every essential map of a metric continuum X onto I^n is weakly confluent.*

Proof. Mazurkiewicz [13] proved this for the case $n = 2$ and said in a footnote that it was true for all n . Howard Cook has pointed out that the same proof works for all n . \square

Lemma 2. *If $\{X_i\}_{i=1}^n$ is a finite family of metric continua and for each i , $f_i: X_i \rightarrow I$ is an onto mapping, then the product mapping*

$$F = \prod_{i=1}^n f_i: \prod_{i=1}^n X_i \rightarrow I^n$$

is essential.

Proof. This is a fairly straightforward application of the Brouwer fixed point theorem; it may well be a standard exercise in dimension theory. Let each X_i be embedded into a copy Q_i of the Hilbert cube. If F is not essential, there is a map $g: \prod_{i=1}^n X_i \rightarrow \dot{I}^n$ which agrees with F on $F^{-1}(\dot{I}^n)$. Then, since \dot{I}^n is an ANR, g extends continuously to a product of closed neighborhoods W_i of X_i in Q_i . Call this extension $G: \prod_{i=1}^n W_i \rightarrow \dot{I}^n$.

Let $\delta > 0$ be such that whenever $x, y \in \prod_{i=1}^n W_i$ and $d(x, y) < \delta$, then $d(G(x), G(y)) < \frac{1}{2}$. Assume, with no loss of generality, that each point of each W_i is at a distance less than δ from some point of X_i . Choose $a_i, b_i \in X_i$ such that $f_i(a_i) = 0$ while $f_i(b_i) = 1$. Let $h_i: I \rightarrow W_i$ be a path such that $h_i(1) = a_i$ and $h_i(0) = b_i$. Then

$$G \circ \prod_{i=1}^n h_i: I^n \rightarrow \dot{I}^n \subseteq I^n$$

is a fixed point free map, a contradiction; and the proof is complete.

A continuum X is *decomposable* provided there exist proper subcontinua $A, B \subseteq X$ such that $X = A \cup B$; otherwise X is *indecomposable*. Equivalently, X is indecomposable if and only if every proper subcontinuum of X is nowhere dense. For a nondegenerate indecomposable continuum X , p and q belong to the same *composant* of X provided some proper subcontinuum W of X contains both p and q . This is

easily seen to be an equivalence relation on X , and for X metric, every equivalence class (composant) is known to be a dense first category F_σ set [12]. If p and q lie in different composants of X , then X is irreducible between them. X is *hereditarily indecomposable* provided every subcontinuum of X is indecomposable. Note that it is immediate from this definition that if X is hereditarily indecomposable and A, B are subcontinua of X with $A \cap B \neq \emptyset$, then either $A \subseteq B$ or $B \subseteq A$.

A metric continuum is *chainable* or *arc-like* if and only if, for each $\varepsilon > 0$, there exists an ε -map $f: X \rightarrow I$. A metric continuum which is both chainable and hereditarily indecomposable is called a *pseudo-arc*. Clearly each nondegenerate subcontinuum of a pseudo-arc is a pseudo-arc. For background material on pseudo-arcs, see for example [2, 3, 4, 5, 6, 7, 10, 11, 15, 16].

The letter P , with or without subscripts, will henceforth denote a fixed pseudo-arc with a given metric ρ .

For any set I , and any space X , the group of product homeomorphisms of X^I will be denoted $\mathcal{G}(X^I)$. A homeomorphism $h \in \mathcal{H}(X^I)$ is *factor preserving* if and only if $h = \theta \circ H$ where $H \in \mathcal{G}(X^I)$ and θ only permutes the factors of X^I . If every self-homeomorphism of X^I is factor-preserving, X^I is *factorwise rigid*.

M is the Menger universal curve. K. Kuperberg, W. Kuperberg, and Transue [8] proved that $M \times M$ is factorwise rigid. The second author [17] extended this result to arbitrary products of M or of Sierpinski universal curves. Lewis [9, problem 60] has asked whether P has the same property. Lysko and Bellamy [3] have given a proof that this is true for the product $P \times P$, using techniques that do not directly generalize to larger products. In the present article, this result is extended to all products of pseudo-arcs.

The next four lemmas summarize the principal facts about pseudo-arcs which will be used here.

Lemma 3 [5] or [16]. P is homogeneous.

Lemma 4 [6]. P has the fixed point property.

Lemma 5 [10]. Given any homeomorphism $h: P \rightarrow P$ and any $\varepsilon > 0$ there is a finite sequence h_1, h_2, \dots, h_n of ε -homeomorphisms of P whose composition $h_1 \circ h_2 \circ \dots \circ h_n$ is h .

Lemma 6. Given any $\varepsilon > 0$ and any p, q which belong to different composants of P , there exists an ε -map $f: P \rightarrow 1$ such that $f^{-1}(0) = \{p\}$ and $f^{-1}(1) = \{q\}$.

Proof. This follows easily from Theorem 1 of [4, p. 44] and the proof thereof. \square

In a product $\prod_{\gamma \in I} X_\gamma$, a set obtained by fixing a single coordinate will be called a *slice*. In particular, the set

$$L(\alpha, p) = \left\{ \langle x_\gamma \rangle_{\gamma \in I} \in \prod_{\gamma \in I} X_\gamma \mid x_\alpha = p \right\}$$

will be called an α -slice, or, if the point p is needed, the p - α slice. The same notation will be used if $y = \langle y_\gamma \rangle_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} X_\gamma$; in this case, $L(\alpha, y) = \{\langle x_\gamma \rangle_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} X_\gamma \mid x_\alpha = y_\alpha\}$. Note that for fixed α , distinct α -slices are disjoint; and if one slice contains another slice, the two are equal. In I^Γ , the 0 - α and 1 - α slices are called the opposite α -faces of I^Γ .

Lemma 7. *A homeomorphism h of a product continuum $\prod_{\gamma \in \Gamma} X_\gamma$ is a product homeomorphism if and only if, for every α , the image under h of every α -slice is an α -slice.*

Proof. Define $h_\alpha: X_\alpha \rightarrow X_\alpha$ by $h_\alpha(x) = y$ where $h(L(\alpha, x)) = L(\alpha, y)$. It is clear that $h = \prod_{\alpha \in \Gamma} h_\alpha$. \square

Henceforth, $\pi_\alpha: \prod_{\gamma \in \Gamma} X_\gamma \rightarrow X_\alpha$ will always denote the projection map, with the exception that, in I^Γ , $\hat{\pi}_\alpha: I^\Gamma \rightarrow I_\alpha = I$ will denote the projection.

Lemma 8. *Let X be a nondegenerate continuum. A homeomorphism $h \in \mathcal{H}(X^\Gamma)$ is factor preserving if and only if the image of every slice under h is a slice.*

Proof. Fix $\alpha \in \Gamma$, and define $A(\beta) = \{p \in X \mid h(L(\alpha, p)) \text{ is a } \beta\text{-slice}\}$. Note that each $A(\beta)$ is open in X . To see this, suppose $x \in A(\beta)$, and $h(L(\alpha, x)) = L(\beta, y)$. For any U open in X such that $y \in U$ but $U \neq X$, there is some open set \hat{V} in $\prod_{\gamma \in \Gamma} X_\gamma$ such that $L(\alpha, x) \subseteq \hat{V}$ and $\pi_\beta \circ h(\hat{V}) \subseteq U$. Then using compactness, there is an open set V in X containing x such that

$$\left\{ \langle z_\gamma \rangle_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} X_\gamma \mid z_\alpha \in V \right\} \subseteq \hat{V}.$$

Thus, if $w \in V$, $\pi_\beta \circ h(L(\alpha, w)) \neq X$, and $h(L(\alpha, w))$ is a β -slice. Thus $x \in V \subseteq A(\beta)$, so that $A(\beta)$ is open. Hence, $\{A(\beta)\}_{\beta \in \Gamma}$ is a covering of X by pairwise disjoint open sets. Since X is connected, $A(\beta) = X$ for some β ; denote this β by $\phi(\alpha)$. Then $\phi: \Gamma \rightarrow \Gamma$ is a function.

To prove that the image of every slice under h^{-1} is a slice, suppose $L(\beta, y)$ is a β -slice. Since $\{h^{-1}(y)\} = \bigcap \{L(\alpha, h^{-1}(y)) \mid \alpha \in \Gamma\}$, it follows that $\{y\} = \bigcap \{h(L(\alpha, h^{-1}(y))) \mid \alpha \in \Gamma\}$. Let \hat{y} be a point differing from y only in the β th coordinate. Then since $\hat{y} \notin \bigcap \{h(L(\alpha, h^{-1}(y))) \mid \alpha \in \Gamma\}$, there is an $\hat{\alpha} \in \Gamma$ such that $\hat{y} \notin h(L(\hat{\alpha}, h^{-1}(y)))$. Then, since $h(L(\hat{\alpha}, h^{-1}(y)))$ is a slice, it must be a β -slice; in particular, $h(L(\hat{\alpha}, h^{-1}(y))) = L(\beta, y)$. Thus, $h^{-1}(L(\beta, y)) = L(\hat{\alpha}, h^{-1}(y))$, as required. Hence, ϕ^{-1} is a function as well, so that $\phi: \Gamma \rightarrow \Gamma$ is bijective. This implies that h is factor-preserving, being a product homeomorphism followed by the permutation ϕ on the factors.

The converse is clear, and the proof is complete. \square

The set function T is defined on the subsets of a continuum X as follows: $T(A)$ is that subset of X such that for $x \in X$, $x \notin T(A)$ if and only if x has a closed,

connected neighborhood W such that $W \cap A = \emptyset$. A substantial body of literature exists on the set function T ; the interested reader is referred to the articles in Sections II and V of [14] and the bibliographies thereof. The next three lemmas give the only facts about T which are needed here.

Lemma 9. *If $\{X_\gamma\}_{\gamma \in \Gamma}$ is a family of indecomposable continua and $L \subseteq \prod_{\gamma \in \Gamma} X_\gamma$ is a slice, then $T(L) = \prod_{\gamma \in \Gamma} X_\gamma$.*

Proof. If W is a subcontinuum of $\prod_{\gamma \in \Gamma} X_\gamma$ and W has nonempty interior, then its projection to every X_γ must be onto. Thus W meets every slice. \square

Lemma 10. *If Γ is nondegenerate and $\prod_{\gamma \in \Gamma} X_\gamma$ is a product of nondegenerate continua and A is a proper subset of a slice in $\prod_{\gamma \in \Gamma} X_\gamma$, then $T(A) \neq \prod_{\gamma \in \Gamma} X_\gamma$.*

Proof. Assume $A \subseteq L(\alpha, p)$ and let $\langle a_\gamma \rangle_{\gamma \in \Gamma} \in L(\alpha, p) - A$. Let U be a closed set with interior in X_α such that $p \notin U$. Then

$$\{\langle x_\gamma \rangle_{\gamma \in \Gamma} \mid x_\alpha \in U \text{ or for every } \gamma \neq \alpha, x_\gamma = a_\gamma\}$$

is a continuum with nonempty interior which misses A . (This is a classical argument of F.B. Jones.) \square

Lemma 11. *If X is a continuum, $A \subseteq X$, and $h: X \rightarrow X$ is a homeomorphism, then $T(h(A)) = h(T(A))$.*

Proof. This is clear. \square

If $B \subseteq \Gamma$ then $Q_B: \prod_{\gamma \in \Gamma} X_\gamma \rightarrow \prod_{\gamma \in B} X_\gamma$ is the projection, $Q_B(\langle x_\gamma \rangle_{\gamma \in \Gamma}) = \langle x_\gamma \rangle_{\gamma \in B}$, with the exception that \hat{Q}_B denotes this projection in I^Γ .

Lemma 12. *Let L be a slice in P^Γ . Suppose $h \in \mathcal{H}(P^\Gamma)$ and that for some α , $\pi_\alpha \circ h(L) \neq P$. Let $f_\gamma: P \rightarrow I$ be an onto map for each $\gamma \in \Gamma$ and assume that $f_\alpha^{-1}(0)$ and $f_\alpha^{-1}(1)$ are singletons lying in different composants of P and missing $\pi_\alpha \circ h(L)$. Let $F = \prod_{\gamma \in \Gamma} f_\gamma: P^\Gamma \rightarrow I^\Gamma$. Then $F(h(L))$ separates I^Γ between its two α -faces. (That is, if C is a continuum in I^Γ which intersects both $\{x \in I^\Gamma \mid x_\alpha = 0\}$ and $\{x \in I^\Gamma \mid x_\alpha = 1\}$, then $F(h(L)) \cap C \neq \emptyset$.)*

Proof. Let $p_0 = f_\alpha^{-1}(0)$ and $p_1 = f_\alpha^{-1}(1)$. Suppose that there is a continuum $A \subseteq I^\Gamma$ with points a_0 and a_1 on the 0- and 1- α -faces, respectively, such that $A \cap F(h(L)) = \emptyset$. Then there is a finite subset B of Γ containing α such that $\hat{Q}_B(A) \cap \hat{Q}_B(F(h(L))) = \emptyset$. Now since $F_B = \prod_{\gamma \in B} f_\gamma: P^B \rightarrow I^B$ is weakly confluent, there is some continuum $K \subseteq P^B$ such that $F_B(K) = \hat{Q}_B(A)$. There are points b_0 and b_1 in K such that $F_B(b_0) = \hat{Q}_B(a_0)$ and $F_B(b_1) = \hat{Q}_B(a_1)$. But then the α th coordinate of b_0 is p_0 and the α th coordinate of b_1 is p_1 . It follows that $\pi_\alpha(Q_B^{-1}(K)) = P_\alpha = P$, since Q_B is

monotone and no proper subcontinuum of P contains $\{p_0, p_1\}$. Suppose U is a closed neighborhood of p_0 in P_α missing $\pi_\alpha \circ h(L)$, and let $M = \{x \in P^\Gamma \mid x_\alpha \in U\}$. Then $M \cup Q_B^{-1}(K)$ is a continuum in P^Γ with nonempty interior which misses $h(L)$. But this can't be, since $T(h(L)) = P^\Gamma$. This contradiction completes the proof. \square

Lemma 13. *Suppose $L \subseteq P^\Gamma$ is a slice and $h \in \mathcal{H}(P^\Gamma)$ and $h(L)$ is not a slice. Then for every $\gamma \in \Gamma$, $\pi_\gamma \circ h(L) = P$.*

Proof. Suppose for some $\alpha \in \Gamma$, $\pi_\alpha(h(L)) \neq P_\alpha$. Let $p_0, p_1 \in P_\alpha - \pi_\alpha(h(L))$, and assume that p_0 and p_1 belong to distinct composants of P_α , and neither belongs to the composant containing $\pi_\alpha(h(L))$. Without loss of generality, assume L is a β -slice, and let $\langle a_\gamma \rangle_{\gamma \in \Gamma} \in L$.

Let $\langle W_k \rangle_{k=1}^\infty$ be a decreasing sequence of nondegenerate subcontinua of P_β whose intersection is $\{a_\beta\}$. Let $W(k) = W_k \times \{\langle a_\gamma \rangle_{\gamma \neq \beta}\}$. By hypothesis, $h(L)$ is not a slice. If $h(L)$ is a proper subset of a slice, then $T(h(L)) \neq P^\Gamma$ by Lemma 10, while by Lemmas 9 and 11, $T(h(L)) = P^\Gamma$. Therefore, $\pi_\alpha(h(L))$ is nondegenerate, and it follows that, for some m , $\pi_\alpha(h(W(m))) \subseteq \pi_\alpha(h(L))$. Choose $b \in W_m$ sufficiently close to a_β that $\pi_\alpha(h(L(\beta, b))) \neq P_\alpha$, and let $J = L(\beta, b)$. Then $\pi_\alpha(h(J)) \cap \pi_\alpha(h(L)) \neq \emptyset$, so that either $\pi_\alpha(h(J)) \subseteq \pi_\alpha(h(L))$ or $\pi_\alpha(h(L)) \subseteq \pi_\alpha(h(J))$. Assume the former; the two cases are identical. There is a finite subset B of Γ , containing α , such that $Q_B(h(L)) \cap Q_B(h(J)) = \emptyset$. Since $Q_B(h(L))$ and $Q_B(h(J))$ are subsets of $\prod_{\gamma \in B} P_\gamma$, there is $\varepsilon > 0$ such that ε is less than the distance from $Q_B(h(J))$ to $Q_B(h(L))$ (with respect to the max metric on $\prod_{\gamma \in B} P_\gamma$). For each $\gamma \in \Gamma$ let $f_\gamma: P \rightarrow I$ be an ε -map, and assume that $f_\alpha^{-1}(0) = p_0$ and $f_\alpha^{-1}(1) = p_1$. Then $F = \prod_{\gamma \in \Gamma} f_\gamma: P^\Gamma \rightarrow I^\Gamma$ is a continuous map, and $F_B = \prod_{\gamma \in B} f_\gamma: P^B \rightarrow I^B$ is an ε -map. Thus, $F_B(Q_B(h(J))) \cap F_B(Q_B(h(L))) = \emptyset$, and since $F_B(Q_B(h(J))) = \hat{Q}_B(F(h(J)))$ and $F_B(Q_B(h(L))) = \hat{Q}_B(F(h(L)))$, it follows that $\hat{Q}_B(F(h(J))) \cap \hat{Q}_B(F(h(L))) = \emptyset$ and that $F(h(J)) \cap F(h(L)) = \emptyset$. Furthermore, by Lemma 12, $F(h(J))$ separates I^Γ between its two α -faces. (Neither $F(h(J))$ nor $F(h(L))$ meets an α -face by composant choices for p_0 and p_1 .) However, $\hat{\pi}_\alpha(F(h(J))) = f_\alpha(\pi_\alpha(h(J)))$, and since $\pi_\alpha(h(J)) \subseteq \pi_\alpha(h(L))$, it follows that $f_\alpha(\pi_\alpha(h(J))) \subseteq f_\alpha(\pi_\alpha(h(L)))$, and thus $\hat{\pi}_\alpha(F(h(J))) \subseteq \hat{\pi}_\alpha(F(h(L)))$. Let $\langle x_\gamma \rangle_{\gamma \in \Gamma}$ and $\langle y_\gamma \rangle_{\gamma \in \Gamma}$ be points in $F(h(L))$ with smallest and largest α th coordinates, respectively. Let

$$A = \{\langle t_\gamma \rangle_{\gamma \in \Gamma} \in I^\Gamma \mid \text{for each } \gamma \neq \alpha, t_\gamma = x_\gamma, \text{ and } 0 \leq t_\alpha \leq x_\alpha\},$$

$$B = \{\langle t_\gamma \rangle_{\gamma \in \Gamma} \in I^\Gamma \mid \text{for each } \gamma \neq \alpha, t_\gamma = y_\gamma, \text{ and } y_\alpha \leq t_\alpha \leq 1\}.$$

Then, each of A and B is a straight line segment meeting $F(h(L))$, and $A \cup B \cup F(h(L))$ is a continuum in I^Γ joining the two α -faces but missing $F(h(J))$. But $F(h(J))$ separates I^Γ between these two faces, and the proof is done by contradiction. \square

Lemma 14. *Suppose $h \in \mathcal{H}(P^\Gamma)$, L is an α -slice and $\pi_\alpha \circ h(L) \neq P_\alpha$. Then $h(L)$ is an α -slice.*

Proof. The proof of Lemma 13 gives this result, too. \square

Lemma 15. Suppose $h \in \mathcal{H}(P^\Gamma)$, $\alpha \in \Gamma$, and there is an α -slice L such that $\pi_\alpha \circ h(L) \neq P_\alpha$. Then for each α -slice L' , $h(L')$ is an α -slice.

Proof. Let $A = \{x \in P \mid h(L(x, \alpha)) \text{ is an } \alpha\text{-slice}\}$, and $B = \{x \in P \mid h(L(x, \alpha)) \text{ is not an } \alpha\text{-slice}\}$. Then $A \neq \emptyset$, since if $z \in L$, $L = L(z_\alpha, \alpha)$ and $h(L)$ is an α -slice. If $B \neq \emptyset$, $A \cup B = P$, and $A \cap B = \emptyset$.

Let $a \in A$ and define x by $h(L(a, \alpha)) = L(x, \alpha)$. Suppose $B \neq \emptyset$ and let $b \in B$. Then $h(L(b, \alpha))$ is not an α -slice, and so by Lemma 13, $\pi_\alpha(h(L(b, \alpha))) = P_\alpha$; in particular, $x \in \pi_\alpha(h(L(b, \alpha)))$, so that $h(L(b, \alpha)) \cap L(x, \alpha) \neq \emptyset$; that is $h(L(b, \alpha)) \cap h(L(a, \alpha)) \neq \emptyset$. Consequently $L(b, \alpha) \cap L(a, \alpha) \neq \emptyset$. Thus $a = b$, which is impossible. Hence, $B = \emptyset$ and $A = P$. \square

Theorem 1. For each finite set B in Γ , let $G(B) = \{h \in \mathcal{H}(P^\Gamma) \mid \text{for each } \alpha \in B, \text{ if } L \text{ is an } \alpha\text{-slice, } h(L) \text{ is an } \alpha\text{-slice}\}$. Then $G(B)$ is an open-closed subgroup of $\mathcal{H}(P^\Gamma)$.

Proof. Clearly $G(B)$ is a group. Also, $G(B) = \bigcap_{\alpha \in B} G(\{\alpha\})$, and so, to show that $G(B)$ is open, all that needs to be shown is that $G(\{\alpha\})$ is open for each α . Fix α . For convenience, let $G(\{\alpha\}) = G$.

Recall that the topology on $\mathcal{H}(P^\Gamma)$ is the compact-open topology. If C is a compact subset of P^Γ and U is open in P^Γ , then $\langle C, U \rangle = \{h \in \mathcal{H}(P^\Gamma) \mid h(C) \subseteq U\}$ is a sub-basic open set in $\mathcal{H}(P^\Gamma)$. To see that some open set containing 1 is in G , suppose $U(\alpha)$ and $V(\alpha)$ are open in P such that $U(\alpha) \cup V(\alpha) = P$, but $\overline{U(\alpha)} \neq P$ and $\overline{V(\alpha)} \neq P$. Then find $H(\alpha)$ and $K(\alpha)$ such that $H(\alpha)$ and $K(\alpha)$ are closed subsets of P such that $H(\alpha) \cup K(\alpha) = P$, $H(\alpha) \subseteq U(\alpha)$ and $K(\alpha) \subseteq V(\alpha)$. Let $U = \{x \in P^\Gamma \mid x_\alpha \in U(\alpha)\}$, $V = \{x \in P^\Gamma \mid x_\alpha \in V(\alpha)\}$, $H = \{x \in P^\Gamma \mid x_\alpha \in H(\alpha)\}$ and $K = \{x \in P^\Gamma \mid x_\alpha \in K(\alpha)\}$. Then $1 \in \langle H, U \rangle \cap \langle K, V \rangle$, which is open in $\mathcal{H}(P^\Gamma)$. But also, $\langle H, U \rangle \cap \langle K, V \rangle \subseteq G$, by Lemma 14. Thus, G is open. But then G is closed, too, for all open subgroups are closed. \square

Theorem 2. Recall that $\mathcal{G}(P^\Gamma)$ denotes the set of product homeomorphisms in $\mathcal{H}(P^\Gamma)$. Then $\mathcal{G}(P^\Gamma) = \bigcap_{\alpha \in \Gamma} G(\{\alpha\})$ and $\mathcal{G}(P^\Gamma)$ is a closed, normal subgroup of $\mathcal{H}(P^\Gamma)$.

Proof. Clearly $\mathcal{G}(P^\Gamma)$ is a closed subgroup of $\mathcal{H}(P^\Gamma)$, since $\mathcal{G}(P^\Gamma) = \bigcap_{\alpha \in \Gamma} G(\{\alpha\})$.

Suppose $h \in \mathcal{G}(P^\Gamma)$ and $g \in \mathcal{H}(P^\Gamma)$. Consider $g \circ h \circ g^{-1}$. Write $h = \prod_{\alpha \in \Gamma} h_\alpha$ where, for $\alpha \in \Gamma$, $h_\alpha \in \mathcal{H}(P)$. For B a finite set of Γ , let $h(B) = \prod_{\alpha \in B} h(B)_\alpha$ where $h(B)_\alpha = 1$ ($\in \mathcal{H}(P)$) for $\alpha \notin B$, and $h(B)_\alpha = h_\alpha$ for $\alpha \in B$. Further, let $h(\tilde{B}) = \prod_{\alpha \in \Gamma} h(\tilde{B})_\alpha$ where $h(\tilde{B})_\alpha = 1$ ($\in \mathcal{H}(P)$) for $\alpha \in B$, and $h(\tilde{B})_\alpha = h_\alpha$ for $\alpha \notin B$.

If $1 \in V$, an open set in $\mathcal{H}(P^\Gamma)$, there is an open set U in $\mathcal{H}(P^\Gamma)$ such that $1 \in U$ and if $f \in U$, then $g \circ f \circ g^{-1} \in V$. (This is because of the continuity of the group operations.) To see that if $\alpha \in \Gamma$, then $g \circ h \circ g^{-1} \in G(\{\alpha\})$, fix α . Since $G(\{\alpha\})$ is open, there is a basic open set U containing 1 such that $g \circ U \circ g^{-1} \subseteq G(\{\alpha\})$. Write

$U = \bigcap_{i=1}^n \langle C(i), U(i) \rangle$, i.e., as an intersection of sub-basic sets. Since $1 \in U$, $C(i) \subseteq U(i)$ for each i . For each i , there are finite collections $\{V(i, 1), \dots, V(i, m(i))\}$ and $\{V'(i, 1), \dots, V'(i, m(i))\}$ of basic open sets of P^Γ such that

$$C(i) \subseteq \bigcup_{j=1}^{m(i)} V(i, j) \subseteq \bigcup_{j=1}^{m(i)} \overline{V'(i, j)} \subseteq \bigcup_{j=1}^{m(i)} V(i, j) \subseteq U(i),$$

and $\overline{V'(i, j)} \subseteq V(i, j)$ for each $j \leq m(i)$. Now $C(i) = \bigcup_{j=1}^{m(i)} (C(i) \cap \overline{V'(i, j)})$ and for $j \leq m(i)$, $D(i, j) = C(i) \cap \overline{V'(i, j)} \subseteq V(i, j)$.

Note that if \mathcal{O} is a basic open set in P^Γ and C is a closed set in \mathcal{O} , then there is a finite set B_i in Γ such that $h(\tilde{B}_i) \in \langle C, \mathcal{O} \rangle$. Thus, there is a finite set B_i in Γ such that

$$h(\tilde{B}_i) \in \bigcap_{j=1}^{m(i)} \langle D(i, j), V(i, j) \rangle \subseteq \langle C(i), U(i) \rangle.$$

Let $B = \bigcup_{i=1}^n B_i$. Then

$$h(\tilde{B}) \in \bigcap_{i=1}^n \left(\bigcap_{j=1}^{m(i)} \langle D(i, j), V(i, j) \rangle \right) \subseteq \bigcap_{i=1}^n \langle C(i), U(i) \rangle = U,$$

and $g \circ h(\tilde{B}) \circ g^{-1} \in G(\{\alpha\})$.

For $\beta \in B$, Lemma 5 can be used to write h_β as a finite composition $h_\beta(1) \circ h_\beta(2) \circ \dots \circ h_\beta(k)$ of homeomorphisms of P such that if $h_\beta(i) = \prod_{\gamma \in \Gamma} f_\gamma$ where $f_\gamma = 1 \in \mathcal{H}(P)$ if $\gamma \neq \beta$, and $f_\beta = h_\beta(i)$, then $g \circ \hat{h}_\beta(i) \circ g^{-1} \in G(\{\alpha\})$ and $g \circ h(\{\beta\}) \circ g^{-1} = (g \circ \hat{h}_\beta(1) \circ g^{-1}) \circ (g \circ \hat{h}_\beta(2) \circ g^{-1}) \circ \dots \circ (g \circ \hat{h}_\beta(k) \circ g^{-1}) \in G(\{\alpha\})$. But then $g \circ h(B) \circ g^{-1} \in G(\{\alpha\})$, since $h(B)$ is a composition of the $h(\{\beta\})$'s for $\beta \in B$ (in any order; they commute) and $h = h(B) \circ h(\tilde{B})$ so that $g \circ h \circ g^{-1} = (g \circ h(B) \circ g^{-1}) \circ (g \circ h(\tilde{B}) \circ g^{-1}) \in G(\{\alpha\})$. This completes the proof. \square

Theorem 3. *Every homeomorphism of a product of pseudo-arcs is factor preserving; that is, products of pseudo-arcs are factorwise rigid.*

Proof. It suffices, by Lemma 8, to prove that the image of every slice under any $h \in \mathcal{H}(P^\Gamma)$ is a slice. Suppose to the contrary that there exists a slice $L \subseteq P^\Gamma$ and an $h \in \mathcal{H}(P^\Gamma)$ such that $h(L)$ is not a slice. By Lemma 13, for every γ , $\pi_\gamma(h(L)) = P$. Let $\langle p_\gamma \rangle_{\gamma \in \Gamma} \in h(L)$, and let $\langle q_\gamma \rangle_{\gamma \in \Gamma}$ be any point in P^Γ differing from $\langle p_\gamma \rangle_{\gamma \in \Gamma}$ in only one coordinate; that is, for some α , $p_\alpha \neq q_\alpha$, but $p_\gamma = q_\gamma$ for $\gamma \neq \alpha$. Let $g_\alpha: P \rightarrow P$ be a homeomorphism such that $g_\alpha(p_\alpha) = q_\alpha$ and let a_α be a fixed point of g_α . For $\gamma \neq \alpha$, let g_γ be the identity map, and $g = \prod_{\gamma \in \Gamma} g_\gamma$ be the product homeomorphism. Since $\pi_\alpha(h(L)) = P$, in particular, $a_\alpha \in \pi_\alpha(h(L))$, and so there exists a point $a = \langle a_\gamma \rangle_{\gamma \in \Gamma} \in h(L)$ such that $\pi_\alpha(a) = a_\alpha$. Since, for each γ , a_γ is a fixed point of g_γ , a is a fixed point of g . Now $g \in \mathcal{G}(P^\Gamma)$, and so $h^{-1} \circ g \circ h \in \mathcal{G}(P^\Gamma)$ also. Consequently, $h^{-1} \circ g \circ h(L)$ is a slice in the same direction as L . However, since $a \in h(L)$, there exists $x \in L$ such that $h(x) = a$. Then $g(h(x)) = g(a) = a$, and $h^{-1} \circ g \circ h(x) = x$. Therefore, $h^{-1} \circ g \circ h(L) \cap L \neq \emptyset$, so that $h^{-1} \circ g \circ h(L) = L$. Thus, since $\langle q_\gamma \rangle_{\gamma \in \Gamma} = g(\langle p_\gamma \rangle_{\gamma \in \Gamma}) \in g(h(L))$; $h^{-1}(\langle q_\gamma \rangle_{\gamma \in \Gamma}) \in h^{-1} \circ g \circ h(L) = L$, and so $\langle q_\gamma \rangle_{\gamma \in \Gamma} \in h(L)$. Thus,

every point of P^I , which differs in only one coordinate from a point of $h(L)$, itself belongs to $h(L)$. However, if B is a finite subset of I , $\langle p_\gamma \rangle_{\gamma \in I} \in h(L)$ and $\langle s_\gamma \rangle_{\gamma \in I} \in P^I$ such that $s_\gamma = p_\gamma$ for each $\gamma \notin B$, there is a finite sequence $\{x(k)\}_{k=0}^n$ of points of P^I such that $x(0) = \langle p_\gamma \rangle_{\gamma \in I}$, $x(n) = \langle s_\gamma \rangle_{\gamma \in I}$, and each $x(k-1)$ differs from $x(k)$ only in one coordinate. Thus if $P' = \{z \in P^I \mid z_\gamma = p_\gamma \text{ for } \gamma \notin B\}$, $P' \subseteq h(L)$. But then $h(L) = P^I$, for $h(L)$ is a closed set. This is a contradiction, since $h(L)$ is a proper subcontinuum of P^I . \square

It is peculiar that homogeneity plays such a strong role in this proof. Is every (finite) product of hereditarily indecomposable continua factorwise rigid?

Acknowledgement

The early stages of this research were done in collaboration with J.M. Lysko, who coauthored the original paper [3] on the subject. Although Lysko declined to appear as coauthor of the present paper, we wish to thank him for several useful conversations early on in the course of the work.

References

- [1] R.F. Arens, A topology for spaces of transformations, *Ann. of Math.* 247 (1946) 480–495.
- [2] D.P. Bellamy, Mapping hereditarily indecomposable continua onto a pseudo-arc, *Lecture Notes in Mathematics* 375, Topology Conference, Virginia Polytechnic Institute and State University, 1973 (Springer, Berlin, 1973) 6–14.
- [3] D.P. Bellamy and J.M. Lysko, Factorwise rigidity of the product of two pseudo-arcs, *Topology Proceedings* 8 (1983) 21–27.
- [4] R.H. Bing, Concerning hereditarily indecomposable continua, *Pacific J. Math.* 1 (1951) 43–51.
- [5] R.H. Bing, A homogeneous indecomposable plane continuum, *Duke Math. J.* 15 (1948) 729–742.
- [6] O.H. Hamilton, A fixed point theorem for pseudo-arcs and certain other metric continua, *Proc. Amer. Math. Soc.* 2 (1951) 173–174.
- [7] B. Knaster, Un continu dont tout sous-continu est indécomposable, *Fund. Math.* 3 (1922) 247–286.
- [8] K. Kuperberg, W. Kuperberg and W.R.R. Transue, On the two homogeneity of Cartesian products, *Fund. Math.* 110 (1980) 131–134.
- [9] I.W. Lewis, Ed., Continua theory problems, *Topology Proceedings* 8 (2) (1983) 361–394.
- [10] I.W. Lewis, Most maps of the pseudo-arc are homeomorphisms, *Proc. Amer. Math. Soc.* 91 (1984) 147–154.
- [11] I.W. Lewis, Stable homeomorphisms of the pseudo-arc, *Canadian J. Math.* 31 (1979) 363–374.
- [12] S. Mazurkiewicz, Un théorème sur les continus indécomposables, *Fund. Math.* I (1920) 35–39.
- [13] S. Mazurkiewicz, Sur l'existence des continus indécomposables, *Fund. Math.* XXV (1935) 327–328.
- [14] L.F. McAuley and M.M. Rao, Eds., *General Topology and Modern Analysis* (Academic Press, New York, 1981).
- [15] E.E. Moise, An indecomposable plane continuum which is homeomorphic to each of its nondegenerate subcontinua, *Trans. Amer. Math. Soc.* 63 (1948) 581–594.
- [16] E.E. Moise, A note on the pseudo-arc, *Trans. Amer. Math. Soc.* 67 (1949) 57–58.
- [17] J.K. Phelps, Homeomorphisms of products of universal curves, *Houston J. Math.* 6 (1980) 127–134.